


1920
H13

C. Hale

Concerning the Occurrence of
Free Pentoses in Plants



LIBRARY
UNIVERSITY OF ILLINOIS
1920

**CONCERNING THE OCCURRENCE OF
FREE PENTOSE IN PLANTS**

BY

CEDRIC HALE

THESIS

FOR THE

DEGREE OF BACHELOR OF SCIENCE

IN

CHEMICAL ENGINEERING

COLLEGE OF LIBERAL ARTS AND SCIENCES

UNIVERSITY OF ILLINOIS

1920

ALICE
H. DEATH
1880

1920
H13

1920

1920

1920

UNIVERSITY OF ILLINOIS

May 25, 1920 191.....

THIS IS TO CERTIFY THAT THE THESIS PREPARED UNDER MY SUPERVISION BY

.....
CEDRIC HALE
.....

ENTITLED.....Concerning the Occurrence of Free Pentoses in Plants.....
.....

IS APPROVED BY ME AS FULFILLING THIS PART OF THE REQUIREMENTS FOR THE

DEGREE OF.....Bachelor of Science.....

.....
in Chemical Engineering
.....

.....
Instructor in Charge

APPROVED :

HEAD OF DEPARTMENT OF.....Chemistry.....

452914

RECEIVED
JAN 10 1900
U.S. DEPT. OF AGRICULTURE

OFFICE OF THE SECRETARY

TO THE SECRETARY OF AGRICULTURE
WASHINGTON, D.C.

FROM THE SECRETARY OF AGRICULTURE
WASHINGTON, D.C.

TO THE SECRETARY OF AGRICULTURE
WASHINGTON, D.C.

FROM THE SECRETARY OF AGRICULTURE
WASHINGTON, D.C.

TO THE SECRETARY OF AGRICULTURE
WASHINGTON, D.C.

FROM THE SECRETARY OF AGRICULTURE
WASHINGTON, D.C.



LIBRARY
100 HILLMAN
000000

ACKNOWLEDGMENT

The author expresses his indebtedness to Dr. Duane T. Englis upon whose suggestion his investigation was undertaken, and under whose direction it was carried out. A sincere appreciation is felt for the encouragement of independent work combined with the feeling that valuable suggestions would be offered when most needed.

VI 11117
1010001101100
58117

Introduction

The purpose of this study is to investigate the effects of the proposed changes on the environment. The study is based on a review of the literature and field observations. The results of the study are presented in the following sections. The first section describes the study area and the proposed changes. The second section discusses the methods used in the study. The third section presents the results of the study. The fourth section discusses the conclusions of the study. The fifth section discusses the implications of the study for future research and policy.

- TABLE OF CONTENTS -

I. Introduction and History	Page 1
II. Theoretical and Experimental	
1. Plan of Experiment	4
2. Preparation of Solutions	4
3. Effect of Refluxing with Ammonium Hydroxide	4
4. Effect of Concentration	6
III. Recovery of Iodide Salts	7
IV. Summary	9
V. Bibliography	10

CONTENTS

1	THE HISTORY OF THE UNITED STATES	1
2	THE HISTORY OF THE UNITED STATES	2
3	THE HISTORY OF THE UNITED STATES	3
4	THE HISTORY OF THE UNITED STATES	4
5	THE HISTORY OF THE UNITED STATES	5
6	THE HISTORY OF THE UNITED STATES	6
7	THE HISTORY OF THE UNITED STATES	7
8	THE HISTORY OF THE UNITED STATES	8
9	THE HISTORY OF THE UNITED STATES	9
10	THE HISTORY OF THE UNITED STATES	10


- CONCERNING THE OCCURRENCE OF FREE PENTOSE IN PLANTS -

I


Introduction and History.

That pentoses occur in plants in the combined state has been a recognized fact since the first study of the existence of sugars in plants. According to Armstrong (1), pentoses are never found as simple sugars but as polysaccharides of high molecular weight and as complex glucosides which are present essentially as skeletal and not as food products; and are, therefore, outside the range of the ordinary plant enzymes. He also points out that the simple sugars more than then complex carbohydrates exert a preponderating influence on pentose formation, the amount of pentoses increasing during the day and decreasing with increase of moisture.

Spoehr (2) brings out the theory of the formation of pentoses from such substances as glucuronic acid. He states that in the disaccharides and polysaccharides, the CH_2OH group is the first to be affected by chemical change with the result that glucuronic acid is formed. Since a general property of these acids is the splitting off of CO_2 from the carboxyl group, glucuronic acid would form l-xylose:



Furthermore, if pentoses are derived from a direct oxidation of hexoses, d-glucose would yield d-arabinose and d-galactose give d-lyxose:




ORIGINAL ARTICLES

THE EFFECT OF THE VARIOUS TYPES OF EXERCISE ON THE
HEART RATE AND BLOOD PRESSURE IN THE ELDERLY
BY DR. J. H. HARRIS, JR., M.D., AND DR. J. H. HARRIS, JR., M.D.
The purpose of this study was to determine the effect of various types of exercise on the heart rate and blood pressure in the elderly. The subjects were divided into three groups: the first group consisted of 10 subjects who were engaged in regular exercise; the second group consisted of 10 subjects who were engaged in irregular exercise; and the third group consisted of 10 subjects who were not engaged in any exercise. The results of the study showed that the heart rate and blood pressure were significantly lower in the first group than in the second and third groups. This suggests that regular exercise has a beneficial effect on the heart and blood pressure in the elderly.

THE EFFECT OF THE VARIOUS TYPES OF EXERCISE ON THE
HEART RATE AND BLOOD PRESSURE IN THE ELDERLY
BY DR. J. H. HARRIS, JR., M.D., AND DR. J. H. HARRIS, JR., M.D.
The purpose of this study was to determine the effect of various types of exercise on the heart rate and blood pressure in the elderly. The subjects were divided into three groups: the first group consisted of 10 subjects who were engaged in regular exercise; the second group consisted of 10 subjects who were engaged in irregular exercise; and the third group consisted of 10 subjects who were not engaged in any exercise. The results of the study showed that the heart rate and blood pressure were significantly lower in the first group than in the second and third groups. This suggests that regular exercise has a beneficial effect on the heart and blood pressure in the elderly.

THE EFFECT OF THE VARIOUS TYPES OF EXERCISE ON THE
HEART RATE AND BLOOD PRESSURE IN THE ELDERLY
BY DR. J. H. HARRIS, JR., M.D., AND DR. J. H. HARRIS, JR., M.D.
The purpose of this study was to determine the effect of various types of exercise on the heart rate and blood pressure in the elderly. The subjects were divided into three groups: the first group consisted of 10 subjects who were engaged in regular exercise; the second group consisted of 10 subjects who were engaged in irregular exercise; and the third group consisted of 10 subjects who were not engaged in any exercise. The results of the study showed that the heart rate and blood pressure were significantly lower in the first group than in the second and third groups. This suggests that regular exercise has a beneficial effect on the heart and blood pressure in the elderly.



It is a fact, however, that in nature d-glucose is almost always found with l-xylose, and that d-galactose is associated with l-arabinose and that the pentose sugars thus found are of the l- and not of the d- series.

It was not until 1914, however, that Davis and Sawyer (3) first brought out the theory that pentoses existed in the free state. They based their conclusions on the fact that a non-fermentable reducing sugar residue was obtained after fermentation with yeast and also that by running the regular Kröber and Tollens determination of pentoses by distillation with hydrochloric acid, a greater precipitate was obtained with phloroglucin than could be accounted for as due to hexoses. They also show that the furfural-like compounds produced from other sugars is of a very small significance in the majority of cases.

However, Davis used a 1% solution of ammonium hydroxide, Sp. Gr..88, for neutralizing the organic acids in the refluxing process of extracting the sugars from the plant material. This seems to leave his work open to criticism on account of the well known isomerizing effect of weak alkalies. This was pointed out as early as 1895 by Lobry de Bruyn (4). Also, the extensive researches of Nef (5) have indicated that a great number of products may be formed by action of alkalies on the monosaccharides in the presence of oxidizing agents and that the nature of the products depends largely on the concentration of those substances. Working with ammonia, he took 24.5 gm. of dextrose in a closed vessel with

200 cc. of 1 equivalent ammonium hydroxide and allowed the solution to stand 74 days. At the end of that time the solution was colored yellow and showed C 4.9 and 85.2% sugar. Also, 10 cc. of the solution needed 28.3 cc. of N/10 HCl for neutralization instead of the 27.4 cc. calculated quantity. The 8% loss of ammonia is probably due to the saturation of the C₆ acids which were formed. Accordingly, ammonium hydroxide must act like traces of caustic alkali upon sugar solutions.

Ruff (6) showed that by using ferric salts with hydrogen peroxide, some oxidation or degradation takes place with the formation of d-arabinose, the ferric salts acting as a catalytic agent.

Therefore, due to Davis' method of treatment with ammonium hydroxide and the use of iron vessels, perhaps enough isomerization of levulose to glucose or some other non-fermentable sugar could have taken place to give the reducing residue. Furthermore, it is probable that some intramolecular change in the sugar or some degradation could have given rise to a product which leads to the production of more furfural than the original hexoses.

1870
The first of the year was a very dry one, and the
winter was unusually early. The first snow fell on
the 1st of November, and the first frost on the 15th.
The weather was very cold, and the ground was
frozen. The first of the year was a very dry one,
and the winter was unusually early. The first snow
fell on the 1st of November, and the first frost on
the 15th. The weather was very cold, and the
ground was frozen. The first of the year was a
very dry one, and the winter was unusually early.

The first of the year was a very dry one, and
the winter was unusually early. The first snow
fell on the 1st of November, and the first frost
on the 15th. The weather was very cold, and
the ground was frozen. The first of the year
was a very dry one, and the winter was
unusually early. The first snow fell on the
1st of November, and the first frost on the
15th. The weather was very cold, and the
ground was frozen. The first of the year was
a very dry one, and the winter was unusually
early. The first snow fell on the 1st of
November, and the first frost on the 15th.
The weather was very cold, and the ground
was frozen. The first of the year was a very
dry one, and the winter was unusually early.

II

Theoretical and Experimental.

The experiment, therefore, was essentially a duplication of Davis' work, using approximately 10 gm. of a mixture of sucrose and invert sugar; endeavoring to ascertain whether any errors could be introduced due to the various influences as mentioned in the introduction to this paper.

Preparation of Solutions.- Twenty grams of pure sucrose was dissolved in 220 cc. of water. 100 cc. of this solution was inverted and made up to 220 cc. (solution A) and 100 cc. was simply made up to 220 cc. (solution B). In this manner, by using 100 cc. of both solutions A and B, approximately 10 gm. of the sugar would be used in a determination.

Refluxing with Ammonium Hydroxide.- The next step consisted in refluxing for eighteen hours, 100 cc. of each solution A and B in 1500 cc. of boiling alcohol to which 5 cc. of concentrated ammonium hydroxide had been added. The resulting solution was then concentrated to 150 cc., made up to 500 cc., cleared with basic lead, and washed almost to 2000 cc. The excess lead was removed with sodium carbonate and the solution made up to 2 liters.

Determination of Pentoses by Distillation - Fifty cc. portions of the above solution was then distilled with HCl according to the regular Krober and Tollen's (7) method of analysis and the precipitate resulting from the addition of phloroglucinol was weighed. The result of this first run showed a phloroglucinol precipitate of .0191 gm. and .0188 gm.

THE HISTORY OF THE
CITY OF BOSTON

From the first settlement of the
English in 1630 to the present time
the city of Boston has been the seat of
the most important and interesting
events in the history of the
New England colonies. It was the
first city in America to be founded
by Englishmen, and it was the
first city in America to be the seat
of a government. It was the first
city in America to be the seat of
a university, and it was the first
city in America to be the seat of
a library. It was the first city in
America to be the seat of a
theater, and it was the first city in
America to be the seat of a
museum. It was the first city in
America to be the seat of a
hospital, and it was the first city in
America to be the seat of a
prison. It was the first city in
America to be the seat of a
court, and it was the first city in
America to be the seat of a
parliament. It was the first city in
America to be the seat of a
university, and it was the first city in
America to be the seat of a
library. It was the first city in
America to be the seat of a
theater, and it was the first city in
America to be the seat of a
museum. It was the first city in
America to be the seat of a
hospital, and it was the first city in
America to be the seat of a
prison. It was the first city in
America to be the seat of a
court, and it was the first city in
America to be the seat of a
parliament.

In carrying out the last distillation, the rate was kept as high as possible (30 cc. every 10 min.) since, according to Van Haarst and Olivier (8), except in the case of l-arabinose, the most complete distillation of the furfural is obtained when the time is short, continued boiling with HCl tending to decompose the furfural.

After the addition of the phloroglucinol, the solution was allowed to stand over night in order to insure complete precipitation. If allowed to stand for a longer time a heavier precipitate is formed which, according to the work of Cunningham and Dorie (9), is w-hydroxymethylfurfuraldehyde. They state that this compound is formed by the action of HCl acid on hexoses, starch, and celluloses, but that owing to its slow formation it does not interfere with the accuracy of the pentose determination as long as aniline acetate is used as an indicator of complete distillation. This compound also differs from the phloroglucid in that the latter is black while the former is brownish-red.

Removal of Hexoses and Determination of Pentoses by Reduction Method.- After the above determination was finished, 1000 cc portions of the 2 liter solution was concentrated in a vacuum to 250 cc. and 50 cc. portions fermented for three to five weeks with *Sach Cereviseae*, cleared with alumina cream, and washed to 100 cc. with boiling water. A regular Browne-Morris copper oxide reduction was run on 50 cc. portions of the solutions.

Results:

cc. $\text{Na}_2\text{S}_2\text{O}_3$	CuO	Arabinose mg.
3.4	.0255	9.2
4.32	.0324	12.0

The next step was a blank Krober-Tollen's distillation on 50cc. portions of a 2 liter solution made by simply diluting 100 cc of each of the two solutions A and B. The results obtained gave a precipitate of .0078 gm. and .0085 gm.

A reduction determination was also made on 50 cc. portions of the above solution, after three to five weeks fermentation, but the results showed no reduction of the copper oxide.

Effect of Concentration.- In the vacuum concentration as carried out above, a slow stream of air was allowed to bubble thru the solution in order to prevent bumping. As this passage of air might result in a slight oxidation to pentose sugars, a distillation was carried out on solutions which had been concentrated in a vacuum without the passage of the air. The results obtained in this way showed a precipitation of .0165 gm. and .0162 gm.

In order to ascertain whether the ammonium hydroxide had an effect on the production of pentoses if allowed to remain in contact with sugar for a shorter time, another determination was made on solutions in which there was no refluxing with the ammonium hydroxide. Here a precipitate was obtained which amounted to .0097 gm. and .0091 gm. Furthermore, the w-hydroxymethylfurfuraldehyde precipitate was very small as compared with the other determinations.

III

Recovery of Iodide Salts.

In the copper oxide reduction method as carried out above and especially in using the Peter's modification method, large quantities of potassium iodide was used. Because of the shortage and cost of this product an attempt was made to recover it from the titration residues.

Stephenson (10) showed that the free iodine could be obtained by treating the solution with nitric acid and hydrochloric acid. Trossian (11) also showed that the iodine could be obtained by treating the residues with hydrogen peroxide. Nothing could be found, however, in regard to the recovery of the iodine in the form of the iodide.

The first step in the recovery consisted in the partial neutralization of the solution containing the sulphates and tartrates of sodium and potassium, the iodide of copper, the sodium thiosulphate, and starch, with sodium hydroxide, leaving the solution only slightly acid in order to prevent the formation of iodates. By passing H_2S into this solution, the copper sulphide is precipitated out, carrying the starch down with it. The solution was filtered, leaving a practically clear solution of the sulphate and iodide salts of sodium and potassium. By evaporating this filtrate these salts crystallize out, giving white crystals of sodium and potassium iodides and sulphates. Since the iodides are soluble in alcohol while the sulphates are not, the former were dissolved in alcohol, best by using a Soxhlet extractor for several hours. The sodium and potassium iodides were then crystallized from the alcoholic

solution.

In order to insure that no free iodine is present, the filtrate, after the CuS has been precipitated out, should be washed with carbon bisulphide in a separatory funnel, the carbon bisulphide dissolving any free iodine.

The above method of recovery gives a sufficiently pure product of sodium and potassium iodides to enable one to use these salts over again in the thiosulphate titrations.

1881. The year of the great earthquake and fire in San Francisco. The city was almost completely destroyed. The population was reduced to about one-third of its former number. The city was in a state of confusion and distress. The people were suffering from the loss of their homes and property. The city was in a state of confusion and distress. The people were suffering from the loss of their homes and property. The city was in a state of confusion and distress. The people were suffering from the loss of their homes and property.

IV

Summary.

1. In a recent paper, pentoses have been reported as occurring in the free state. The evidence for their presence is based on the facts (1) that after fermentation of the hexoses with brewer's yeast a non-fermentable residue remains and (2) more furfural is produced by distillation of a portion of the original solution with dilute hydrochloric acid than could be accounted for from the hexose alone. The extraction of the sugars involved refluxing with ammoniacal alcohol and extensive concentrations.
2. It is shown in the present paper that a reproduction of this process of extraction and concentration with pure sucrose and invert sugar gives results comparable to those reported. Hence, the apparent free pentoses are probably derived from the hexoses.
3. In the absence of ammonia some furfural is obtained, but no residue, non-fermentable with brewer's yeast.
4. The formation of the apparent pentoses is due primarily to the action of air in contact with the ammoniacal solution of the sugars.


The first of these is the fact that the
the second is the fact that the
the third is the fact that the
the fourth is the fact that the
the fifth is the fact that the
the sixth is the fact that the
the seventh is the fact that the
the eighth is the fact that the
the ninth is the fact that the
the tenth is the fact that the
the eleventh is the fact that the
the twelfth is the fact that the
the thirteenth is the fact that the
the fourteenth is the fact that the
the fifteenth is the fact that the
the sixteenth is the fact that the
the seventeenth is the fact that the
the eighteenth is the fact that the
the nineteenth is the fact that the
the twentieth is the fact that the
the twenty-first is the fact that the
the twenty-second is the fact that the
the twenty-third is the fact that the
the twenty-fourth is the fact that the
the twenty-fifth is the fact that the
the twenty-sixth is the fact that the
the twenty-seventh is the fact that the
the twenty-eighth is the fact that the
the twenty-ninth is the fact that the
the thirtieth is the fact that the
the thirty-first is the fact that the
the thirty-second is the fact that the
the thirty-third is the fact that the
the thirty-fourth is the fact that the
the thirty-fifth is the fact that the
the thirty-sixth is the fact that the
the thirty-seventh is the fact that the
the thirty-eighth is the fact that the
the thirty-ninth is the fact that the
the fortieth is the fact that the
the forty-first is the fact that the
the forty-second is the fact that the
the forty-third is the fact that the
the forty-fourth is the fact that the
the forty-fifth is the fact that the
the forty-sixth is the fact that the
the forty-seventh is the fact that the
the forty-eighth is the fact that the
the forty-ninth is the fact that the
the fiftieth is the fact that the
the fifty-first is the fact that the
the fifty-second is the fact that the
the fifty-third is the fact that the
the fifty-fourth is the fact that the
the fifty-fifth is the fact that the
the fifty-sixth is the fact that the
the fifty-seventh is the fact that the
the fifty-eighth is the fact that the
the fifty-ninth is the fact that the
the sixtieth is the fact that the
the sixty-first is the fact that the
the sixty-second is the fact that the
the sixty-third is the fact that the
the sixty-fourth is the fact that the
the sixty-fifth is the fact that the
the sixty-sixth is the fact that the
the sixty-seventh is the fact that the
the sixty-eighth is the fact that the
the sixty-ninth is the fact that the
the seventieth is the fact that the
the seventy-first is the fact that the
the seventy-second is the fact that the
the seventy-third is the fact that the
the seventy-fourth is the fact that the
the seventy-fifth is the fact that the
the seventy-sixth is the fact that the
the seventy-seventh is the fact that the
the seventy-eighth is the fact that the
the seventy-ninth is the fact that the
the eightieth is the fact that the
the eighty-first is the fact that the
the eighty-second is the fact that the
the eighty-third is the fact that the
the eighty-fourth is the fact that the
the eighty-fifth is the fact that the
the eighty-sixth is the fact that the
the eighty-seventh is the fact that the
the eighty-eighth is the fact that the
the eighty-ninth is the fact that the
the ninetieth is the fact that the
the ninety-first is the fact that the
the ninety-second is the fact that the
the ninety-third is the fact that the
the ninety-fourth is the fact that the
the ninety-fifth is the fact that the
the ninety-sixth is the fact that the
the ninety-seventh is the fact that the
the ninety-eighth is the fact that the
the ninety-ninth is the fact that the
the hundredth is the fact that the

V
Bibliography.

- I "The Simple Carbohydrates and the Glucosides". (1912)
- 2 "The Carbohydrate Economy of Cacti". (1919)
- 3 J. of Agr. Sc. 6, 406, (1914)
- 4 Rec. des trav. chim. des Pays-Bas, 14, 203-216, (1895)
- 5 Annalen, 403, 204, (1913)
- 6 Ber. chem. Ges., 31, 1573, (1898)
- 7 U. S. Chemistry Bureau Bulletin #107.
- 8 Chem. Weekblad, 11, 918.
- 9 Bio-chem. J. 8, 438.
- 10 Analyst, 43, 165-6, (1918).
- 11 J. Ind. Eng. Chem. 6, 83, (1914)

Y 55610
CH 1159 26 27 28 29 30
10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30

UNIVERSITY OF ILLINOIS-URBANA



3 0112 082200392